

## **Erythromelalgia- a clinical sign never to be missed**

Ovallath Sujith<sup>1</sup>, Dsouza Michelle<sup>2</sup>

<sup>1</sup>DM (Neurology), Fellowship in Movement Disorder (Canada), HOD

<sup>2</sup>MBBS, Junior Resident

Department of Neurology, Kannur Medical College, Anjarakandy, Kannur, Kerala- 670612

### **\*Corresponding author**

Dr. Sujith Ovallath

Email: [sujithok@gmail.com](mailto:sujithok@gmail.com)

**Abstract:** Erythromelalgia is a rare clinical condition due to post ganglionic sympathetic dysfunction. Diagnosis of this disorder and differentiating it from other common burning pain syndromes like Raynaud's phenomenon and small fibre neuropathy is often difficult. We report a patient who presented with redness, pain and burning sensation of hands, who was managed successfully with aspirin.

**Keywords:** Thrombocythemia, Erythromelalgia, SCN9A gene, Aspirin

### **INTRODUCTION:**

Erythromelalgia is a rare disease affecting mainly the peripheral extremities, manifested by pain and burning sensation of hands and feet.[1] The term was coined by Silas Weir Mitchell in 1878 and is derived from 3 Greek words 'erythros' (red), 'melos' (extremity), and 'algos' (pain). Primary (idiopathic) and secondary (associated with an underlying disease) forms have been described. The present report is about a middle aged man presented with thrombocytomic erythromelalgia, a rare form of autonomic dysfunction which was treated effectively by antiplatelet agent.


### **CASE HISTORY:**

A 58yrs old gentleman was evaluated for pain and burning sensation of bilateral hands especially at the finger tips of 2 yrs duration. He is a manual labourer and persistent pain interfered with his day today activities. Pain and burning sensation increases on exposure to heat and during activities and reduced on dipping the hands in the cold water. No convincing evidence of any other medical illness in the past or any significant family history was obtained. Patient had several hospital admissions without much relief. Examination revealed moderate built with normal vitals and normal joints. Local examination of bilateral hands revealed slight bluish discoloration with mottling appearance [Figure 1.].




**Fig-1:Bluish discoloration with mottling appearance in affected limbs.**

Systemic examinations were otherwise within normal limits. On routine lab investigations revealed normal parameters except for an elevated platelet count: 6.3 lac/cumm. His previous records showed platelet count 9.4lac/cumm. Repeated blood examination confirmed the presence of thrombocythemia. Peripheral smear showed normocytic normochromic anaemia, neutrophilic leucocytosis, thrombocytosis [Figure 2.]



**Fig-2: Peripheral smear showing thrombocytosis**

Nerve conduction study was normal, rheumatoid factor was positive initially but on repeated examination after a few weeks it became negative, X-ray chest did not reveal any evidence of cervical rib. HIV test was negative, MRI cervical spine showed mild disc bulges without any neural element compression [Figure 3.].



**Fig-3: MRI cervical spine showing mild disc bulges without any neural element compression**

Protein electrophoresis showed normal composition of albumin and globulin fractions. No monoclonal band was seen. ANA was negative. Initial clinical diagnosis was Raynaud's phenomenon and the patient was started on Nifedepine which worsened the pain. We reviewed the diagnosis and, on retrospective questioning he reported worsening of pain when hand is immersed in hot water. In view of thrombocytopenia and pain the diagnosis of erythromelalgia was considered and the patient was started on Aspirin 325 mg BID. With one dosage of Aspirin the patient improved markedly. He became completely symptoms free within few days. He is being continued with Aspirin 50mg BD with no recurrence of symptoms.

## DISCUSSION:

In this case the clinical symptoms, presence of thrombocytopenia and improvement on Aspirin suggest the clinical diagnosis of erythromelalgia. Erythromelalgia also known as Michell's disease, was first described by Sir Silas Weir Mitchell in 1878[2]. It is a rare disease which affects the peripheral extremities mostly involving hands and feet. It usually evident in adulthood[2]. But there are many childhood[3] and infantile[4] cases of erythromelalgia reported in the literature. Many classification of Erythromelalgia have been proposed. Recently on basis of clinical, laboratory, and histopathological studies, it has been classified into three distinct types- erythromelalgia in thrombocytopenia, primary erythromelalgia, and secondary erythromelalgia[5]. But sometimes thrombocytopenic erythromelalgia has been grouped under secondary erythromelalgia[6].

The pathophysiology of erythromelalgia depends on the cause. Primary erythromelalgia is inherited in autosomal dominant manner. The root cause is mutation in the human gene SCN9A, which encodes for alpha subunit of voltage gated sodium channels called Na (v) [7]. These voltage gated sodium channels mainly present on postganglionic sympathetic nervous system and c-fibres of dorsal root ganglion. Normally postganglionic sympathetic system controls the cutaneous vascular tone and response to body cooling.[8] In primary Erythromelalgia, mutated voltage gated sodium channels present in the post ganglionic sympathetic system alters the responses to various stimuli and results in manifestations. The nociceptors and warm receptors in human skin are innervated by c fibres.[9] In primary erythromelalgia mutated voltage gated sodium channels present in c-fibres causes hyper excitability of c-fibres which results in manifestations of erythromelalgia.

Secondary erythromelalgia is due to underlying systemic diseases like SLE, myeloproliferative diseases, polycythemia Vera, connective tissue disorders, neuropathies, spinal cord diseases, carcinoma of the colon and thyroid, and astrocytomas[10]. The symptoms of erythromelalgia may precede the onset of a myeloproliferative disease by a median of 2 1/2 years [6]. Thrombocytopenic erythromelalgia is due to increased platelet counts resulting in platelet thrombi and hence causes arteriolar occlusion and fibrosis[11]. Manifestations of erythromelalgia is a triad of redness, pain and burning sensation of peripheral extremities[12]. The symptoms aggravate on exposure to heat and hanging the affected extremities downward. Exposure to cold and elevation of affected limbs can decrease the symptoms[2]. The cases of erythromelalgia has to be differentiated from Raynaud's phenomenon, the differentiating points are shown in Figure 4.

Erythromelalgia	Raynaud's disease
1. Sympathetic dysfunction	Sympathetic overactivity
2. Hyperemic	Hypoperfusion
3. Extremities are red and warm	Blue and cold
4. Symptoms are relieved by cold	Symptoms aggravate on exposure to cold
5. Aggravated by calcium channel blockers like nifedipine	Relieved by calcium channel blockers

**Fig- 4: Differences between Raynaud's phenomenon and Erythromelalgia**

In rare cases gangrene and ulceration may occur. On clinical examination the affected part appears to be red, dusky and mottled with warmth. The other clinical findings may give a clue regarding the underlying systemic diseases.

Supportive evidences such total blood count and histopathological studies may give a great clue regarding the underlying diseases and helps in classifying the type of disease and in excluding other diagnoses[13]. Infrared thermography can also help in diagnosis of conflicted cases. Thermography of an erythromelalgic patient may reveal increased temperature in the affected area. Hence detecting and visualizing the hyper thermal area can assist in diagnosing the patient, assessing the therapeutic results, and understanding the disease course of erythromelalgia[12].

There is no definitive treatment for erythromelalgia[14]. Aspirin (100 -300mg/day) is the treatment of choice specially in thrombocytomic erythromelalgia as in our case[15]. Aspirin inhibits platelet aggregation hence results in dramatic relief of symptoms [16]. Positive response may also helps in confirming the diagnosis. Other oral medications like sodium channel blockers –lidocaine, mexiletine have been tried with lesser effect[7].Surgical measures like sympathectomy have been experimented in cases which are resistant to all medical measures[17].

Following up the patient is mandatory as in our case for serial tests of blood counts. As we stated earlier erythromelalgia may precedes the onset of myeloproliferative diseases. Hence blood counts have to be monitored for early detection and bone marrow examination in suspected cases.

## CONCLUSION

Patients presenting with isolated features of burning sensation and redness of peripheral extremities without much of other clinical findings, the diagnosis of

erythromelalgia has to be considered. Raynaud's phenomenon and small fibre neuropathy are other close differentials. Therapeutic trial of Aspirin is very useful in arriving at a clinical diagnosis of erythromelalgia.

## REFERENCES:

1. Albuquerque LG, Franca ER, Kozmhinsky V, Querino MC, Morais AG; Primary erythromelalgia: case report. *Anais brasileiros de dermatologia*. 2011; 86(1):131-4.
2. Norton JV, Zager E, Grady JF; Erythromelalgia: diagnosis and classification. *The Journal of foot and ankle surgery: official publication of the American College of Foot and Ankle Surgeons*. 1999; 38(3):238-41.
3. Al-Minshawy SM, El-Mazary AA; An Egyptian child with erythromelalgia responding to a new line of treatment: a case report and review of the literature. *Journal of medical case reports*. 2014; 8:69.
4. Ahogo KC, Menet V, Modiano P, Lasek A; [Inconsolable crying revealing primary erythromelalgia in a 6-month-old infant]. *Archives de pediatrie: organe officiel de la Societe francaise de pediatrie*. 2014; 21(3):299-301.
5. Drenth JP, van Genderen PJ, Michiels JJ; Thrombocytomic erythromelalgia, primary erythromelalgia, and secondary erythromelalgia: three distinct clinicopathologic entities. *Angiology*. 1994; 45(6):451-3.
6. Kurzrock R, Cohen PR; Erythromelalgia and myeloproliferative disorders. *Archives of internal medicine*. 1989; 149(1):105-9.
7. Wu MT, Huang PY, Yen CT, Chen CC, Lee MJ; A novel SCN9A mutation responsible for primary erythromelalgia and is resistant to the treatment of sodium channel blockers. *PloS one*. 2013; 8(1):e55212.
8. Strom NA, Meuchel LW, Mundy DW, Sawyer JR, Roberts SK, Kingsley-Berg SM, *et al.*; Cutaneous sympathetic neural responses to body cooling in type 2 diabetes mellitus. *Autonomic neuroscience: basic & clinical*. 2011; 159(1-2):15-9.
9. Hallin RG, Torebjork HE, Wiesenfeld Z; Nociceptors and warm receptors innervated by C fibres in human skin. *Journal of neurology, neurosurgery, and psychiatry*. 1982; 45(4):313-9.
10. Ghosh SK, Bandyopadhyay D, Ghoshal L; Painful red hands and feet. *Cleveland Clinic journal of medicine*. 2012; 79(5):356-8.
11. Van Genderen PJ, Lucas IS, van Strik R, Vuzevski VD, Prins FJ, van Vliet HH, *et al.*; Erythromelalgia in essential thrombocythemia is characterized by platelet activation and endothelial cell damage but not by thrombin generation. *Thrombosis and haemostasis*. 1996; 76(3):333-8.

12. Kang BC, Nam da J, Ahn EK, Yoon DM, Cho JG; Secondary erythromelalgia - a case report. *The Korean journal of pain*. 2013; 26(3):299-302.
13. Davis MD, O'Fallon WM, Rogers RS, 3rd, Rooke TW; Natural history of erythromelalgia: presentation and outcome in 168 patients. *Archives of dermatology*. 2000; 136(3):330-6.
14. Ljubojevic S, Lipozencic J, Pustisek N. Erythromelalgia; *Acta dermatovenerologica Croatica: ADC*. 2004; 12(2):99-105.
15. Barbui T; What is the standard treatment in essential thrombocythemia. *International journal of hematology*. 2002; 76 Suppl 2:311-7.
16. Bertele V, Tomasiak M, Falanga A, Cerletti C, de Gaetano G; Aspirin inhibits platelet aggregation but not because it prevents thromboxane synthesis. *Lancet (London, England)*. 1982; 2(8301):775.
17. Shiga T, Sakamoto A, Koizumi K, Ogawa R; Endoscopic thoracic sympathectomy for primary erythromelalgia in the upper extremities. *Anesthesia and analgesia*. 1999; 88(4):865-6.